



Fact Sheet

United States Nuclear Regulatory Commission
Office of Public Affairs
Washington DC 20555
Telephone: 301/415-8200 E-mail: opa@nrc.gov

Biological Effects of Radiation

Radiation

Radiation is all around us. It is naturally present in our environment and has been since the birth of this planet. Consequently, life has evolved in an environment which has significant levels of ionizing radiation. It comes from outer space (cosmic), the ground (terrestrial), and even from within our own bodies. It is present in the air we breathe, the food we eat, the water we drink, and in the construction materials used to build our homes. Certain foods such as bananas and brazil nuts naturally contain higher levels of radiation than other foods. Brick and stone homes have higher natural radiation levels than homes made of other building materials such as wood. Our nation's capitol, which is largely constructed of granite, contains higher levels of natural radiation than most homes.


Levels of natural or background radiation can vary greatly from one location to the next. For example, people residing in Colorado are exposed to more natural radiation than residents of the east or west coast because Colorado has more cosmic radiation at a higher altitude and more terrestrial radiation from soils enriched in naturally occurring uranium. Furthermore, a lot of our natural exposure is due to radon, a gas from the earth's crust that is present in the air we breathe.

The average annual radiation exposure from natural sources to an individual in the United States is about 300 millirem (3 millisieverts). Radon gas accounts for two-thirds of this exposure, while cosmic, terrestrial, and internal radiation account for the remainder. No adverse health effects have been discerned from doses arising from these levels of natural radiation exposure.

In addition, man-made sources of radiation from medical, commercial, and industrial activities contribute another 60 mrem (0.6 mSv) to our annual radiation exposure. One of the largest of these sources of exposure is medical x-rays. Diagnostic medical procedures account for about 40 mrem (0.5 mSv) each year. In addition, some consumer products such as tobacco, fertilizer, welding rods, gas mantles, luminous watch dials, and smoke detectors contribute another 10 mrem (0.1 mSv) to our annual radiation exposure.

A typical breakdown between natural background radiation and artificial sources of radiation is shown in the pie chart below. It shows natural radiation contributes about 82% of the annual dose to the population while medical procedures contribute most of the remaining 18%. Both natural and artificial radiations affect us in the same way.

Sources of Radiation Exposure



Above background levels of radiation exposure, the NRC requires that its licensees limit maximum radiation exposure to individual members of the public to 100 mrem (1mSv) per year, and limit occupational radiation exposure to adults working with radioactive material to 5,000 mrem (50 mSv) per year. NRC regulations and radiation exposure limits (contained in Title 10 of the Code of Federal Regulations under Part 20) are consistent with recommendations of national and international scientific organizations and with practices in other developed nations.

Biological Effects of Radiation

We tend to think of biological effects of radiation in terms of their effect on living cells. For low levels of radiation exposure, the biological effects are so small they may not be detected. The body has defense mechanisms against damage induced by radiation as well as by chemical carcinogens. Consequently, biological effects of radiation on living cells may result in three outcomes: (1) injured or damaged cells repair themselves, resulting in no residual damage; (2) cells die, much like millions of body cells do every day, being replaced through normal biological processes; or (3) cells incorrectly repair themselves resulting in a biophysical change.

The associations between radiation exposure and the development of cancer are mostly based on populations exposed to relatively high levels of ionizing radiation (e.g., Japanese atomic bomb survivors, and recipients of selected diagnostic or therapeutic medical procedures). Cancers associated with high dose exposure include leukemia, breast, bladder, colon, liver, lung, esophagus, ovarian, multiple myeloma, and stomach cancers. Department of Health and Human Services literature also suggests a possible association between ionizing radiation exposure and prostate, nasal cavity/sinuses, pharyngeal and laryngeal, and pancreatic cancer. The effects of radiation can be seen clearly only at doses much higher than are allowed by Federal regulations.

The period of time between radiation exposure and the detection of cancer is known as the latent period. Those cancers that may develop as a result of radiation exposure are indistinguishable from those that occur naturally or as a result of exposure to other chemical carcinogens. Furthermore, National Cancer Institute literature indicates that other chemical and physical hazards and lifestyle factors (e.g., smoking, alcohol consumption, and diet) significantly contribute to many of these same diseases.

Although radiation is known to cause cancers at high doses and high dose rates, currently there are no data to unequivocally establish the occurrence of cancer following exposure to low doses and dose rates -- below about 10,000 mrem (9100 mSv). Studies of a population exposed to chronic low-levels of radiation above normal background have shown no adverse biological effects. This population includes occupationally exposed radiation workers and people living in areas having high levels of background radiation -- above 1,000 mrem (10 mSv) per year-- such as Denver, Colorado.

Even so, the radiation protection community conservatively assumes that any amount of radiation may pose some risk for causing cancer and hereditary effect, and that the risk is higher for higher radiation exposures. A linear, no-threshold (LNT) dose response relationship is used to describe the relationship between radiation dose and the occurrence of cancer. This dose-response model suggests that any increase in dose, no matter how small, results in an incremental increase in risk. The LNT hypothesis is accepted by the NRC as a conservative model for determining radiation dose standards recognizing that the model may over estimate radiation risk.

High radiation doses tend to kill cells, while low doses tend to damage or alter the genetic code (DNA) of irradiated cells. High doses can kill so many cells that tissues and organs are damaged immediately. This in turn may cause a rapid body response often called Acute Radiation Syndrome. The higher the radiation dose, the sooner the effects of radiation will appear, and the higher the probability of death. This syndrome was observed in many atomic bomb survivors in 1945 and emergency workers responding to the 1986 Chernobyl nuclear power plant accident. Approximately 131 plant workers and firefighters battling the fire at the Chernobyl power plant received high radiation doses (70,000 to 1,340,000 mrem 9700 to 13,400 mSv) and suffered from acute radiation sickness. Of these, 31 died from their radiation injuries. Fortunately, 100 of these firefighters and plant workers recovered from their immediate illnesses.

Because radiation affects different people in different ways, it is not possible to indicate what dose is needed to be fatal. However, it is believed that 50% of a population would die within thirty days after receiving a dose, over a period ranging from a few minutes to a few hours, between 325,000 to 450,000 mrem (3250 to 4500 mSv). This would vary depending on the health of the individuals before the exposure and the medical care received after the exposure. These doses expose the whole body to radiation in a very short period of time (minutes to hours). Similar exposure of only parts of the body will likely lead to more localized effects, such as skin burns.

Conversely, low doses--less than 20,000 mrem (200 mSv)-- spread out over long periods of time (years to decades) don't cause an immediate problem to any body organ. The effects of low doses of radiation, if any, also occur at the level of the cell, and thus changes may not be observed for many years (usually 5-20 years) after exposure.

Genetic effects and the development of cancer are the primary health concerns attributed to radiation exposure. The likelihood of cancer occurring after radiation exposure is about five times greater than a genetic effect. The latter includes increased still births, congenital abnormalities, infant mortality, childhood mortality, decreased birth weight, and altered sex ratio. Genetic effects are the result of a mutation produced in the reproductive cells of an exposed individual that are passed on to their offspring. These effects may appear in the exposed person's direct offspring, or may appear several generations later, depending on whether the altered genes are dominant or recessive. None of the mutated creatures in the science fiction literature and cinema have been observed in humans. Although radiation-induced genetic effects have been observed in laboratory animals (given very high

doses of radiation), no statistically significant increase in genetic effects has been observed among the children born to atomic bomb survivors from Hiroshima and Nagasaki.

December 2000